


Progress in IR8 matching for β^* squeeze

Y.PAPAPHILIPPOU

thanks to R. deMaria, S. Fartoukh, M. Giovannozzi, A. Lombardi T. Risselada

IR8 squeeze history


- Since V6.5, IR8 phase advance at injection changed to accommodate phase advance difference in IR3 and IR7 (O. Bruning, LTC 31/03/04)
 - From $(\mu_x, \mu_y) = (3.204, 2.804)$ to $(3.183, 2.974)$ for beam1 and $(3.059, 2.782)$ for beam2
- Strength files for optics with β^* between 10 and 1m computed by A.Lombardi and T.Risselada
 - Not a smooth transition in quad strengths
- Using V6.500 optics, R. deMaria achieved unsqueezed optics for beam 2 and squeezed for beam1 up to $\beta^* = 3.4\text{m}$ (LOC, 06/12/05)
 - Not the correct phase advance between MKI and TDI, TCLI
 - Large β -function near Q6
 - Strength of MQX not optimised ($>205\text{T/m}$)




IR8 squeeze strategy

- Using optics V6.500 (unequal phase advances for beam1 and beam2)
- Use R. de Maria's “Jacobian” matching algorithm
- First find a $\beta^*=10\text{m}$ optics with the correct total phase advance and setting MQX values to 205 T/m
- From this point start a squeeze until algorithm fails
- Constraint always β near Q6 to values below 350-400m (as at injection)
- Check the aperture and reiterate if needed, adding more constraints
- Compute the crossing scheme and bumps (“nominal” with horizontal crossing angle and “fancy” with crossing in both planes) and check again the aperture
- Proceed for beam1 and for beam2, independently


Squeeze beam1



Squeeze beam1



Squeeze beam1



Quad strengths

